Using SNAP® Test Kits*

Whole blood/serum/plasma sample test procedure



1. Dispense 3 drops of sample and 4 drops of conjugate (5 drops for cardiac tests) into a disposable sample tube.



2. Gently invert the sample tube 4-5 times to mix.



3. Pour the entire contents of the sample tube into the sample

well of a SNAP® device.

Fecal sample test procedure



1. Swab sample and place swab tip into tube. Bend bulb to break seal and release conjugate.



2. Squeeze and release bulb 3 times to mix sample and conjugate.



3. Squeeze bulb to dispense 5 drops into the sample well of a SNAP® device.



4. When color first appears in the activation circle, press firmly to activate. You will hear a distinct "snap."





5. When the appropriate development time has passed, read the result.


BLUE DOT=POSITIVE

Any color development in the sample spot indicates a positive result.

The SNAP® cPL™ Test, SNAP® fPL™ Test and SNAP® Feline proBNP Test are the exceptions because they provide comparative results.

Interpreting results

Infectious Disease Tests




SNAP® 4Dx® Plus—Screen dogs annually for six vector-borne diseases.



SNAP® **Feline Triple**®—Test all new to household, sick and at-risk cats and cats with respiratory or vomiting symptoms.



SNAP® Parvo — Test dogs that present with sudden onset of lethargy, vomiting, fever and diarrhea.



SNAP® Heartworm RT—Screen annually for canine heartworm.



SNAP® FIV/FeLV Combo — Test all new to household, at-risk and symptomatic cats.



SNAP® *Giardia*—Test dogs and cats that present with diarrhea.

KEY



(8) Development time in minutes.

Store at room temperature or in refrigerator.

Store in refrigerator.

ALL components MUST be at room temperature before running the test.

Tech Tip

To help remember how many drops you need, count the number of letters in the color of the liquid:

R-E-D 3 drops of blood, serum or plasma B-L-U-E 4 drops of blue conjugate **G-R-E-E-N 5** drops of green conjugate

B-R-O-W-N 5 drops of fecal mixture

*Before beginning any test, refer to the package insert for complete instructions.



SNAP® **Lepto**—Test dogs when leptospirosis is suspected.

Pancreatic Tests



Abnormal Sample spot is the same color or darker than reference spot.



Normal Sample spot is lighter than reference spot.

SNAP® cPL™—Test dogs that present with vomiting, anorexia and/or abdominal pain.

SNAP® fPL™ —Test cats that present with lethargy, decreased appetite, dehydration, weight loss, vomiting and/or diarrhea.

Cardiac Tests



Abnormal Sample spot is the same color or darker than reference spot.



Normal Sample spot is lighter than reference spot.





